



CPSC 424

Differential Geometry of Curves

© Alla Sheffer & Wolfgang Heidrich



Syllabus

Curves in 2D and 3D

- Implicit vs. Explicit vs. Parametric curves
- Parametric Curves: Basis functions and Splines
- Splines: Bezier and Hermite
- Other bases
- **Elementary differential geometry**
- Subdivision Curves
- 2D Implicits: Analytical and Neural


Surfaces

© Alla Sheffer & Wolfgang Heidrich

Motivation

Parameterization:

- Curve representations we have looked at have fixed parameterization
 - Bézier
 - B-Spline
 - Rational
- Indicates fixed speed for point traveling along the curve



Want:

- Change speed without affecting shape

Motivation

Continuity between curves

- So far included parameterizations
- Example:
 - Bézier continuity constraints depend on relative interval lengths
 - B-Spline continuity depends on interval lengths in knot sequence

Want:

- Geometric concept for continuity that ignores parameterization

Motivation

Differential Geometry helps to:

- Understand parameterizations
- Understand impact of parameterizations on derivatives
 - *Tangent vector, curvature, ...*
- Change parameterization as required
- Derive geometric definition of continuity

For our purposes:

- Look at 3D curves
- 2D as special case

Regularity

Definition:

- Differentiable parametric curve $F(t) : [a, b] \mapsto \mathcal{R}^3$ is called regular if

$$F'(t) \neq 0, \forall t \in [a, b]$$

- (I.e. if the tangent vector is not 0 anywhere)

Note:

- Bézier curves not necessarily regular...

Equivalence / Reparameterization

Definition

- Two regular curves

$$F(t) : [a, b] \mapsto \mathcal{R}^3 \quad G(t) : [c, d] \mapsto \mathcal{R}^3$$


are geometrically equivalent $F \cong G$ if there is a strictly monotonic, differentiable function

$$\varphi(t) : [a, b] \mapsto [c, d]$$

with

$$F(t) = G(\varphi(t))$$

Equivalence / Reparameterization



Question

Are the curves

$F(t) = (t, t)$ t in $[0, 1]$ and $G(t) = (2t, 2t)$ t in $[0, 0.5]$

geometrically equivalent?

- A. Yes
- B. No
- C. Not enough information

Arc Length

Definition

- Arc length of regular curve $F(t) : [a, b] \mapsto \mathcal{R}^3$ given as

$$s(t) := \int_a^t \|F'(\tilde{t})\| d\tilde{t}$$

Parameterization by arc length

$$G(s) \text{ with } G(s(t)) = F(t)$$

- Note: this is a **canonical** representation for any curve
- Point is traveling along G with constant speed 1

Arc Length Example

$$F(t) = (t, t) \quad t \text{ in } [0, 1]$$

$$F'(t) = ? \quad G(s) = ? \quad s(t) := \int_a^t \|F'(\tilde{t})\| d\tilde{t}$$

$$F(t) = (\sin(t), \cos(t)) \quad t \text{ in } [0, \pi]$$

$$F'(t) = ? \quad G(s) = ?$$

$$G(s) \text{ with } G(s(t)) = F(t)$$

Curvature

Definition

- Let G be a curve parameterized by arc length
- We introduce the following terms:
 - Unit tangent $T(s) := G'(s)$
 - Curvature vector $K(s) := G''(s)$
 - Curvature $\kappa(s) := \|K(s)\|$
 - Principal normal $N(s) := K(s) / \kappa(s)$
 - Binormal $B(s) := T(s) \times N(s)$



General Setting

- To translate into general parameterization use derivation rules

— e.g. Unit tangent $T(t) := \frac{dG}{ds} = \frac{dF}{dt} \frac{dt}{ds} = \frac{F'}{\|F'\|}$



Curvature

- What is Curvature?


Curvature

- General parameterization (2D)

$$k(t) = \frac{x'(t)y''(t) - y'(t)x''(t)}{\left(x'(t)^2 + y'(t)^2\right)^{3/2}}$$

Corresponds to radius of osculating circle $R=1/k$

Measures curve bending



© Alla Sheffer & Wolfgang Heidrich

Frenet Frame

Theorem:

- Curvature vector and tangent vector are perpendicular:

$$K(s) \perp T(s)$$

Note:

- Therefore, T, N, and B form an orthonormal coordinate frame
- This is called the Frenet Frame

<https://www.desmos.com/3d/ei5yo6m9al>

© Alla Sheffer & Wolfgang Heidrich



Torsion

With a similar argument to curvature we get

$$B'(s) = \tau(s) \cdot N(s)$$

Note:

- B' is the torsion vector
- τ is the torsion, and indicates how much the curve twists out of the plane ($\tau=0$ means perfectly planar)

<https://www.desmos.com/3d/fd2zkrdg8w>

© Alla Sheffer & Wolfgang Heidrich



Fundamental Theorem of Curves

Theorem:

- For given functions $\kappa(s)$, $\tau(s)$ there exists exactly one (except for rotations and translations) unique curve that is parameterized by arc length and has curvature $\kappa(s)$, and torsion $\tau(s)$

Proof:

- Quite complex, see for example
 - do Carmo
Differential Geometry of Curves and Surfaces

© Alla Sheffer & Wolfgang Heidrich



Geometric Continuity

Definition:

- Two curves

$$F_1(t) : [a, b] \mapsto \mathcal{R}^3, F_2(t) : [b, c] \mapsto \mathcal{R}^3$$

are G^k-continuous (geometrically continuous of degree k), if there are reparameterizations

$$G_1(t) \cong F_1(t) \text{ and } G_2(t) \cong F_2(t)$$

that are C^k continuous, i.e.:

$$G_1^l(t) = G_2^l(t), l = 0 \dots k$$

at shared parameter interval endpoint

© Alla Sheffer & Wolfgang Heidrich



Geometric Continuity

Note:

- In practice, if two curves are G^k continuous, then there always is a reparameterization for F_2 that is C^k continuous to F_1
 - I.e. only one of the two curves needs to be reparameterized*
- If we reparameterize both curves, we can normalize the tangent vector at the transition point to unit length
 - Local arc length parameterization*

© Alla Sheffer & Wolfgang Heidrich